

3/PRTS

1

10/500891
DT04 Rec'd PCT/PTO 07 JUL 2004

PHOSPHINIC PSEUDOPEPTIDE DERIVATIVES FOR THE SELECTIVE
INHIBITION OF THE C-TERMINAL ACTIVE SITE OF ANGIOTENSIN
I CONVERTING ENZYME (ACE)

5

DESCRIPTION

Technical field

10 The present invention relates to the use of phosphinic
pseudopeptide derivatives for the preparation of a
medicinal product for selectively inhibiting the
C-terminal active site of human angiotensin I
converting enzyme (ACE), i.e. without affecting the
N-terminal active site of ACE.

15

Such a medicinal product may be used in the prevention
and treatment of various cardiovascular pathologies in
man.


20 The present invention also relates to novel phosphinic
pseudopeptide derivatives, to pharmaceutical composi-
tions containing them and to processes for preparing
the said phosphinic pseudopeptide derivatives.

25 **Prior art**

Angiotensin I converting enzyme (ACE) is a central
component in the regulation of arterial pressure and in
the homeostasis of the various physiological functions
30 of cardiovascular tissue. These actions appear to
depend partly on:

- the maturation of a powerful vasoconstrictor, angiotensin II, via cleavage of the C-terminal end of angiotensin I, the inactive peptide, with ACE, and
- the degradation via ACE of a powerful vasodilator, bradykinin.

These actions are illustrated below.



10

Arterial hypertension, and also cardiac tissue diseases, result from deregulation of the various vasoconstrictive hormones. Re-establishing the equilibrium between vasoconstrictors and vasodilators, for the benefit of the latter, is one of the main therapeutic objectives of the medicinal products used in human clinical treatment to remedy arterial hypertension and cardiac tissue diseases. It is thus understood how the inhibition of ACE can participate towards these objectives, by preventing the formation of angiotensin II and potentiating bradykinin. ACE inhibitors are used in human clinical treatment not only to reduce

20

arterial hypertension, but also to preserve the functions of cardiac tissue, as described in references [1] and [2].

5 The cloning of ACE followed by the determination of its primary structure showed, surprisingly, the presence of two active sites in this enzyme, as described in reference [3]. Via site-directed mutagenesis, it has been possible to prove that these two active sites are
10 fully functional, i.e. capable of cleaving physiological substrates of ACE, such as angiotensin I and bradykinin (references [4] and [5]).

Despite all the studies performed for more than 20
15 years on ACE, it is still not known whether the presence of two active sites in mammalian ACE, resulting from duplication of an ancestral gene, corresponds to a particular functional role. However, the recent discovery that, *in vivo*, in man, the peptide
20 Ac-SDKP (N-acetyl Ser-Asp-Lys-Pro) is essentially cleaved via the N-terminal active site of ACE, argues in favour of a distinct functional role for each of the active sites of ACE (reference [6]).

25 These considerations have led researchers to attempt to develop inhibitors capable of highly selectively discriminating between the two active sites of ACE, in order to furnish tools capable of establishing *in vivo* the functional role of each of the sites of ACE.

In this regard, it is important to underline that all the inhibitors used to date in clinical studies are mixed ACE inhibitors, i.e. inhibitors that simultaneously block the two active sites of ACE.

5

The first inhibitor that selectively blocks the N-terminal site of ACE, RXP407, which is a phosphinic pseudopeptide, has recently been developed (references [7] and [8]). This inhibitor, which is not metabolized
10 in rats and mice, is, moreover, capable of inhibiting the degradation of the peptide N-acetyl Ser-Asp-Lys-Pro (Ac-SDKP) *in vivo* in mice (reference [9]). Thus, the injection of RXP407, by blocking the N-terminal site of ACE, would prevent the *in vivo* degradation of Ac-SDKP.


15

This inhibitor forms the subject of a preclinical study in the said animal, which is directed towards demonstrating its usefulness for protecting haematopoietic tissue during chemotherapy treatment.

20

However, no inhibitor capable of discriminating between the two active sites of ACE by interacting essentially with the C-terminal site of ACE has been described to date. However, it would be desirable to have available
25 such inhibitors. The reason for this is that, besides their value in experimental and clinical research, they would have the major advantage, over standard mixed ACE inhibitors, of not interfering with the physiological functions associated with the activity of the
30 N-terminal site of ACE, for instance the metabolism of the peptide Ac-SDKP.

It has been demonstrated that phosphinic peptides represent a generic family of compounds capable of very strongly inhibiting zinc metallopeptidases, the peptidase family to which ACE belongs, as may be seen in references [9] to [16]. In these compounds, the role of the PO_2^- group is to interact with the zinc atom located in the active site of these enzymes.



Besides the presence of the PO_2^- group, the chemical nature of the residues P2, P1, P1' and P2' plays a deciding role in ensuring the selectivity of the interactions between a particular phosphinic peptide and a given zinc metallopeptidase (see references [8], [12] and [13]). Thus, the presence of quite specific residues in positions P2, P1, P1' and P2' makes it possible to obtain selective inhibitors, which inhibit only certain zinc metallopeptidases. Such selectivity may be an essential factor in the context of an *in vivo*

use of these inhibitors. Specifically, it is estimated that the *in vivo* toxicity of certain inhibitors is predominantly due to their lack of selectivity for a given target.

5

According to these principles, the search for inhibitors capable of selectively blocking the C-terminal site of ACE consisted in identifying, in the family of phosphinic compounds, particular residues located in positions P1, P1' and P2', which give the inhibitors the capacity to selectively interact with the C-terminal site of ACE.

10

Description of the invention


15

This research has led to the discovery that the presence of a pseudoproline residue in phosphinic pseudopeptides constitutes an essential element for obtaining selective inhibitors of the C-terminal site of ACE.

20


Thus, one subject of the present invention is the use of at least one phosphinic pseudopeptide derivative comprising the amino acid sequence of formula (I) below:

25



in which:

- 5 - R_2 and R_3 , which are identical or different, represent the side chain of a natural or unnatural amino acid, the sequence:



10

also possibly forming the Pro (proline) residue, and

- R_5 represents a hydrogen atom, a pharmacologically acceptable counterion, or a group capable of forming an *in vivo* hydrolysable phosphinic ester;
- 15

for the manufacture of a medicinal product capable of selectively inhibiting the C-terminal site of angiotensin I converting enzyme.

20

In this sequence, the PO_2 group may be in the form PO_2^- by being associated with a hydrogen atom or a pharmacologically acceptable counterion, for example K^+ , Na^+ ,








NH_4^+ or any other pharmacologically acceptable metal or non-metal ion. The nature of the counterion is irrelevant since, in water, the charged groups are dissociated.

5

The PO_2 group may also be in the form of phosphinic ester hydrolysable *in vivo*. In this case, the pseudopeptide is of the prodrug type, and, after *in vivo* hydrolysis of the ester, it generates the active
10 form of the pseudopeptide.

Groups of this type that may be used for R_5 are described in particular in reference [20].


15 Examples of such groups that may be mentioned are groups corresponding to the following formulae:



In these formulae, t-Bu represents the tert-butyl group.


5

According to one particular embodiment of the invention, the phosphinic pseudopeptide derivative corresponds to formula (II) below:



in which:

- 5 - R_1 represents a protecting group for an amine function, or an amino acid or a peptide protected with a protecting group for an amine function,
- R_2 and R_3 , which may be identical or different,
- 10 represent the side chain of a natural or unnatural amino acid, the sequence:



15 also possibly forming the Pro residue,

- R_4 represents a hydrogen atom or a pharmacologically acceptable counterion, and
- 20 - R_5 is as defined above.

In the formulae given above, R_2 and R_3 represent the side chain of a natural or unnatural amino acid, for example a pseudoamino acid.

5 The natural amino acids may be chosen from alanine, arginine, asparagine, aspartic acid, cysteine, glutamine, glutamic acid, glycine, histidine, isoleucine, leucine, norleucine, lysine, methionine, phenylalanine, proline, hydroxyproline, serine, threonine, tryptophan,
10 tyrosine, valine, nitrophenylalanine, homoarginine, thiazolidine and dehydroproline.

A pseudoamino acid may be defined as an amino acid in which the amino or carbonyl function has been replaced
15 with another chemical group.

In formula (II) mentioned above, the group R_1 may be a common protecting group for an amine function in peptide chemistry. Such protecting groups are well
20 known to those skilled in the art and are illustrated, for example, in the book entitled "Protective Groups in Organic Synthesis", Second Edition, T.W. Greene and P.G.M. Wuts, John Wiley & Sons, Inc., pages 309-315 [17]. As examples of such groups that may be used in
25 the invention, mention may be made of acetyl and benzyloxycarbonyl groups.

R_1 may also represent a natural or unnatural amino acid or a peptide whose terminal amine function is protected
30 with a common protecting group such as those described above.

According to the invention, as will be seen herein-
below, the presence of the pseudoproline residue is
essential to obtain the selectivity towards the
5 C-terminal active site of ACE, but the nature of the
side chains present in R_2 and R_3 also plays an
important role in the selectivity of the interactions
of the derivatives used according to the invention with
the N-terminal and C-terminal sites of ACE.


10

Good results as regards the inhibition of the
C-terminal site of ACE have been obtained with pseudo-
peptides in which the group R_2 represents the benzyl,
methyl or phenylethyl group, i.e. the side chain of
15 phenylalanine, alanine and homophenylalanine.

For R_3 , good results have been obtained when R_3
represents the side chain of alanine, arginine or
tryptophan, or when the sequence $-\text{NH}-\text{CH}(\text{R}_3)-\text{CO}-$
20 represents a Pro residue.

Generally, R_4 and R_5 represent a hydrogen atom.

According to one preferred embodiment of the invention,
25 the phosphinic pseudopeptide derivative corresponds to
the following formula:



(pseudo-peptide G)

The phosphinic pseudopeptide derivatives that may be used in accordance with the invention have been found to be capable of selectively inhibiting the C-terminal active site of angiotensin I converting enzyme, and thus of controlling *in vivo* the physiological levels of angiotensin II - which plays a central role in controlling arterial pressure and homeostasis of the cardiovascular functions in man - without, however, interfering with the metabolism of bradykinin or with that of the peptide Ac-SDKP.

Their use as active principles in a medicinal product is thus liable to find numerous applications in the prevention and treatment of human cardiovascular pathologies, and especially pathologies in which bradykinin is thought to take little part, for instance the prevention of atherosclerosis.


Among the phosphinic pseudopeptide derivatives whose use is envisaged according to the invention, some of them have never been described in the literature.

Thus, a subject of the invention is also a phosphinic pseudopeptide derivative comprising the amino acid sequence of formula (I) mentioned above, in which:

5


- R_2 represents the side chain of a natural or unnatural amino acid,
- the sequence:

10



forms the Pro residue:

15



and


- R_5 represents a hydrogen atom, a pharmacologically acceptable counterion or a group capable of forming an *in vivo* hydrolysable phosphinic ester.

20


Among these derivatives, the ones that are especially preferred are those corresponding to the formula (II) mentioned above, in which:

25

- R_1 represents a protecting group for an amine function, or an amino acid or a peptide protected with a protecting group for an amine function,
- 5 - R_2 represents the side chain of a natural or unnatural amino acid,
- the sequence:




forms the Pro residue:



- R_4 represents a hydrogen atom or a pharmacologically acceptable counterion,
 - R_5 represents a hydrogen atom, a pharmacologically acceptable counterion or a group capable of forming an *in vivo* hydrolysable phosphinic ester.
- 20

More particularly, the preferred phosphinic pseudo-peptide derivative is the one of formula:




(pseudo-peptide G)

A subject of the invention is also a pharmaceutical composition comprising at least one phosphinic pseudo-peptide derivative corresponding to formula (II) mentioned above, in which:


- R_2 represents the side chain of a natural or unnatural amino acid,

10

- the sequence:




15 forms the Pro residue:



and

- R₅ represents a hydrogen atom, a pharmacologically acceptable counterion or a group capable of forming an *in vivo* hydrolysable phosphinic ester.

In this composition, the phosphinic pseudopeptide derivative preferably corresponds to the formula:




(pseudo-peptide G)

With regard to the text hereinabove, such a pharmaceutical composition is especially liable to find numerous applications in the prevention and treatment of various cardiovascular pathologies in man.


The phosphinic pseudopeptide derivatives corresponding to formula (II) mentioned above, in which R₄ and R₅ represent a hydrogen atom, may be prepared via a process comprising the following steps:

- 1) reacting a compound of formula (III):




in which R_1 and R_2 are as defined above, with the compound of formula (IV):

5




in which Ac represents the acetyl group and Et represents the ethyl group, to obtain the compound of formula (V):


10




2) converting compound (V) into compound (VI) by
 15 reacting compound (V) with sodium borohydride:




3) protecting the hydroxyl group of compound (VI) with
an adamantyl group Ad to give the compound of
5 formula (VII):



4) saponifying compound (VII) to give the compound of
10 formula (VIII):



5) coupling the compound of formula (VIII) with the
15 amino acid of formula (IX) or (X):



in which R_3 is as defined above, and

- 5 6) removing the protecting group Ad.

According to this process, the phosphinic block of formula (VIII) comprising the pseudoproline is first synthesized, and peptide coupling of this phosphinic
10 block with the desired amino acid is then performed.

Advantageously, the peptide coupling step 5) is performed by solid-phase peptide synthesis using as solid phase a resin substituted with the amino acid of
15 formula (IX) or (X), the N-terminal end of which will have been protected beforehand with an Fmoc (fluorenylmethoxycarbonyl) group.

If necessary, the phosphinic function of the pseudo-peptide of formula (II) in which R_5 represents a
20 hydrogen atom may then be esterified or salified, by reacting it with suitable reagents.

The esterification may be obtained by coupling with an alcohol of formula R_5OH in which R_5 represents a group
25 capable of forming an *in vivo* hydrolysable phosphinic ester, for example using the process described in reference [20] (method A).

The esterification may also be performed by reaction with a halide of formula R_5X in which R_5 represents a group capable of forming an *in vivo* hydrolysable phosphinic ester. This reaction may be performed under alkaline conditions using the process described in reference [20] (method B).

Before performing this esterification, the carboxylic acid function(s) of the pseudopeptide is (are) protected with suitable protecting groups, which are subsequently removed using standard techniques.

When it is desired to salify the phosphinic function of the pseudopeptide of formula (II) in which R_5 is a hydrogen atom, to replace this hydrogen atom with a pharmaceutically acceptable counterion, the pseudopeptide is reacted with a suitable base containing this counterion, for example NaOH, KOH or NH_4OH .

The same technique may be used to salify the terminal carboxylic group of the pseudopeptide, in order to replace the hydrogen atom with a pharmaceutically acceptable counterion.

Other characteristics of the invention will emerge more clearly on reading the rest of the description that follows, which relates to examples for preparing pseudopeptide derivatives in accordance with the invention and for demonstrating their properties, with reference to the attached drawings.

Needless to say, this description hereinbelow is given for illustrative purposes and with no limitation of the subject of the invention.

5

Brief description of the drawings

Figure 1 shows the synthesis of synthons that are useful for the preparation of the phosphinic pseudo-peptides in accordance with the invention.

10

Figure 2 shows the chromatogram obtained during the purification of the pseudopeptide G by high performance liquid chromatography (HPLC).

15

Figure 3 shows the *in vivo* effects of pseudopeptide G on the cleavage of angiotensin I to angiotensin II.

20

Figure 4 shows the *in vivo* effects of pseudopeptide G on the cleavage of bradykinin.

Figure 5 shows the *in vivo* effects of pseudopeptide G and of RPX407 (selective inhibitor of the N-terminal site of ACE) on the cleavage of the peptide Ac-SDKP.


25

Detailed description of the embodiments

The synthesis of the phosphinic pseudopeptides was performed according to the synthetic scheme described in Figure 1.

30

This figure shows the steps of the process resulting in the synthons of the formula (VIII):



5

in which:

- R_1 represents the benzyloxycarbonyl (Cbz) group, and
- 10 - R_2 is the phenyl group (compounds 1a, 3a, 4a, 5a and 6a), the phenylethyl group (compounds 1b, 3b, 4b, 5b and 6b) or the methyl group (compounds 1c, 3c, 4c, 5c and 6c).

15 **Example 1: Preparation of compound 6a**

1) Preparation of compound 1a

20 This aminophosphinic acid derivative is prepared according to the procedure described by Baylis [18], and the enantiomer of R configuration is then obtained via recrystallization according to the protocol reported by Baylis [18].

2) Preparation of compound 2a

This compound is obtained according to the procedure published by Villieras et al. [19]. The product
5 obtained was characterized by NMR:

^1H -NMR (250 MHz, CDCl_3): 7.03 (t, 1H), 5.93 (m, 1H),
4.1 (m, 2H), 2.6 (m, 1H), 2.34 (m, 2H), 1.95 (s, 3H),
1.82 (m, 1H), 1.18 (t, 3H).

10 3) Preparation of compound 3a

A mixture of compound 1a (3.2 g, 10 mmol) and of hexamethyldisilazane (10.5 mL, 50 mmol) under a flow of argon is heated at 110°C for 3 hours. Compound 2
15 (5.5 g, 12 mmol) is added at this temperature and this solution is stirred for 4 hours at 90°C. This solution is cooled to 70°C, 10 mL of absolute ethanol EtOH are added dropwise thereto and the mixture is stirred at 70°C for 30 minutes. After evaporating off the
20 solvents, the residue is dissolved in 5% NaHCO_3 (10 mL) and 5 mL of hexane. After 3 extractions with ethyl acetate EtOAc (3 × 5 mL), the crude product is obtained after evaporating off the solvent. Purification on a column of silica, using a chloroform/methanol/acetic
25 acid mixture (7/0.3/0.3) as mobile phase, gives 4 g of pure compound 3a, in the form of a white solid (89% yield).

The NMR characterization of this product is based on
30 COSY, TOCSY and HMQC experiments:

¹H-NMR (250 MHz, CDCl₃): 1.27 (t, ³J_{HH} = 7.1 Hz, 3H, CH₂CH₃), 1.95-3.00 (m, 5H, PCH(CH₂)₂, PhCHH), 3.13-3.45 (m, 2H, PCH, PhCHH), 4.02-4.31 (m, 2H, CH₂CH₃), 4.34-4.63 (m, 1H, PCH), 4.78-5.05 (m, 2H, OCH₂Ph), 5.71 (d, 1H, NH, ³J_{HH} = 11.3 Hz, I), 5.78 (d, 1H, NH, ³J_{HH} = 10.8 Hz, II), 6.91-6.99 (d, 1H, C=CH), I/II), 7.02-7.34 (m, 10H, aryl).

¹³C-NMR (62 MHz, CDCl₃): 14.2/14.2 (CH₂-CH₃), 26.2/26.5 (PCHCH₂), 32.7/32.8 (PCHCH₂CH₂), 34.5 (CH₂Ph), 41.8 (d, ¹J_{PC} = 87.7 Hz, PCHC, I), 43.1 (d, ¹J_{PC} = 87.3 Hz, PCHC, II), 50.5 (d, ¹J_{PC} = 99.7 Hz, PCHN, I), 50.5 (d, ¹J_{PC} = 100.5 Hz, PCHN, II), 61.1/61.2 (CH₂CH₃), 66.8 (OCH₂Ph), 126.6, 127.7, 127.8, 127.9, 127.9, 128.4, 128.4, 129.4, 132.2, 132.3, 132.9, 136.5, 136.6, 137.1, 137.3, (aryls), 148.1 (d, ²J_{PC} = 8.7 Hz, =CCO, I), 148.1 (d, ²J_{PC} = 9 Hz, =CCO, II), 156.1 (d, ²J_{PC} = 5.5 Hz, CONH, I), 156.2 (d, ²J_{PC} = 5.7 Hz, CONH, II), 165.3 (d, ²J_{PC} = 2.7 Hz, COOEt).

³¹P-NMR (100 MHz, CDCl₃): 46.89, 48.13.

The indications I and II correspond to the different diastereoisomers.

Elemental analysis:

Theoretical values:

C: 61.80%, H: 6.27%, N: 3.00%.

Experimental values:

C: 61.89%, H: 6.23%, N: 2.98%

4) Preparation of compound 4a

Compound 3a (1.4 g, 3.06 mmol) and $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ (1.09 g, 9.2 mmol) are dissolved in a mixture of THF (12.4 mL)/
5 methanol (7.7 mL). NaBH_4 (0.58 g, 15.4 mmol) is added portionwise to this solution over 30 minutes, at -30°C . This mixture is stirred for a further 10 minutes at -30°C . The solvents are evaporated off and the product is extracted into a mixture of EtOAc (25 mL) and 1N HCl
10 (20 mL, pH 1).

The organic phase is collected and washed with water (10 mL) and then dried with Na_2SO_4 . After evaporating off the solvents, the product is purified on a column
15 of silica with a chloroform/methanol/acetic acid (7/0.3/0.3) mobile phase. 1.28 g of compound 4a are obtained (91% yield).

The analysis by negative-mode mass spectrometry (mass
20 observed $\text{MH}^- = 458.48$, expected mass = 459.47) is in accordance with the chemical structure of compound 4a.

Elemental analysis:

25 Theoretical values:

C: 61.78%, H: 6.63%, N: 3.00%

Experimental values:

C: 61.98%, H: 6.31%, N: 3.08%

5) Preparation of compound 5a

The adamantylation of compound 4a is performed according to the protocol described by Yiotakis et al. [14].

1-Adamantyl bromide (538 mg, 2.5 mmol) and Ag₂O (577 mg), divided into 5 equal portions, are added over 1 hour to a solution of compound 4a (1.03 g, 2.24 mmol) in chloroform. After 2 hours, 0.5 eq of AdBr and 0.5 eq of Ag₂O are added and the mixture is refluxed for 10 hours. After evaporating off the solvents, the crude product is purified on a column of silica, using a chloroform/isopropanol mixture (9.8/0.2) as mobile phase. Compound 5a is obtained in pure form in a yield of 96% (1.27 g).

Analysis by mass spectrometry, positive mode: mass observed MH⁺ = 594.21, expected mass = 593.1.

20

Elemental analysis:

Theoretical values:

C: 67.76%, H: 7.53%, N: 2.32%

25

Experimental values:

C: 67.49%, H: 7.58%, N: 2.24%

6) Preparation of compound 6a

After diluting compound 5a (1.1 g, 1.85 mmol) in methanol (20 mL), 2 mL of 4N NaOH are added. After
5 stirring for 6 hours, TLC monitoring of the reaction confirms the complete saponification of the starting material. After evaporating off the solvent, the product is taken up in a mixture of water (10 mL) followed by addition of EtOAc (15 mL) and acidification
10 to pH 1 with 1N HCl. The residue is taken up into the organic phase and the extraction procedure is repeated twice. The combined organic phases are dried over Na₂SO₄ and the solvents are then evaporated off. The pure compound 6a is obtained in a yield of 94% (0.98
15 g).

Analysis by mass spectrometry, positive mode: mass observed MH⁺ = 566.15, expected mass = 565.26.

Elemental analysis:

20

Theoretical values:

C: 67.95%, H: 7.13%, N: 2.48%

Experimental values:

25 C: 67.64%, H: 7.30%, N: 2.40%

Example 2: Preparation of compound 6b

The same procedure as in Example 1 is followed to
30 prepare compound 6b, starting with compound 1b, which

is prepared according to the procedure described in [18].


The analysis by mass spectrometry of compound 6b gave the following results: expected mass = 579.27, mass observed MH^+ = 580.29.

Example 3: Preparation of compound 6c

10 The same procedure as in Example 1 is followed to prepare compound 6c, starting with compound 1c, which is prepared according to the procedure described in [18].

15 Analysis by mass spectrometry of compound 6c gave the following results: expected mass = 489.23, observed mass MH^+ = 490.11.

Example 4: Preparation of the pseudopeptide G of formula:



(pseudo-peptide G)


This pseudopeptide was synthesized on a solid phase using a standard protocol of solid-phase peptide synthesis. Wang resin substituted with an Fmoc-Trp (732 mg, 0.58 mmol) is suspended in N-methylpyrrolidone (NMP) (5 mL) and stirred for 5 minutes. After removal of the NMP by filtration, 10 mL of piperidine at a concentration of 20% in NMP are added and the mixture is stirred for 15 minutes. After filtration, the resin is washed with the following solvents: NMP (7×10 mL), CH₂Cl₂ (3×10 mL) and Et₂O (2×10 mL). 2 mL of NMP, diisopropylethylamine DIEA (749 mg, 5.76 mmol) and compound 6a (360 mg, 0.64 mmol) diluted in NMP (2 mL) and 2-(1H)benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate HBTU (730 mg, 1.92 mmol, diluted in 3 mL of NMP) are then added to the reactor. The mixture is stirred for 24 hours. After filtration, the resin is washed with NMP (4×7 mL) and CH₂Cl₂ (5×7 mL). A trifluoroacetic acid TFA/CH₂Cl₂/H₂O/triisopropylsilane (90/7.5/1.25/1.25) solution is then added to the reactor and the mixture is stirred for 3 hours (deprotection phase). After filtration, the filtrate containing pseudopeptide G is recovered, the solvent is evaporated off and the product is dissolved in H₂O. After freeze-drying, pseudopeptide G is purified by reverse-phase HPLC (Vydac column, C18, semi-preparative).

Figure 2 shows the chromatogram obtained. In this figure, 4 peaks corresponding to the 4 diastereoisomers present in pseudopeptide G are observed (these 4 peaks


have an identical mass spectrum, mass observed $MH^+ = 618.23$, expected mass = 617.23). Only peak 1 shows inhibitory power with respect to ACE.

- 5 The NMR characterization of pseudopeptide G, peak 1 HPLC, is based on COSY, TOCSY and HMQC experiments:
 1H -NMR (250 MHz, DMSO) 4.93 (d, 2H, CH_2-O-Ph), 4.01 (m, 1H, $CH-CH_2-Ph$), 2.77 (m, 1H, $CH-CH_2-Ph$), 3.08 (m, 1H, $CH-CH_2-Ph$), 3.09 (α -pseudo-Proline), 1.77 (β -pseudo-Proline),
10 Proline), 1.56 (γ -pseudo-Proline), 1.79 (δ -pseudo-Proline), 2.49 (ϵ -pseudo-Proline), 4.50 (α -Trp), 3.13 (β, β -Trp), 7.6 NH, 8.3 NH, aryls 7.35-6.9.


- 15 **Examples 5 to 7: Preparation of pseudopeptides B, C and D of formulae:**



pseudo-peptide B



pseudo-peptide C



pseudo-peptide D

5

Pseudopeptides B, C and D were synthesized on a solid phase with Wang resins substituted with alanine (B), proline (C) and arginine (D), using the protocol described for the preparation of pseudopeptide G. HPLC purification of these pseudopeptides made it possible to isolate diastereoisomers of these pseudopeptides capable of inhibiting ACE.

Analysis of the pseudopeptides by mass spectrometry confirms the structure of these pseudopeptides.


Pseudopeptide B: expected mass 502.19; observed mass 503.21.

Pseudopeptide C: expected mass 528.53; observed mass 529.11.


5 Pseudopeptide D: expected mass 587.25; observed mass 587.24.

Examples 8 and 9: Preparation of pseudopeptides E and F of formulae:

10



pseudo-peptide E



pseudo-peptide F

Pseudopeptides E and F are obtained via solid-phase synthesis, starting with compounds 6b and 6c and following the protocol described for the preparation of pseudopeptide G. After purification by reverse-phase
5 C18 HPLC, the first fraction collected for each of these pseudopeptides was found to be capable of inhibiting ACE.

Analysis by mass spectrometry gave the following
10 results:

Pseudopeptide E: expected mass 541.20; observed mass 542.26.

Pseudopeptide F: expected mass 631.24; observed mass 632.26.

15

Example 10: Determination of the inhibition constants of pseudopeptides A to G with respect to the N-terminal and C-terminal sites of ACE

20 Human recombinant ACE is used for this determination. Curves of inhibition of ACE with pseudopeptides A to G are obtained using the quenched-fluorescence substrate Mca-Ala: Mca-Ala-Ser-Asp-Lys-DpaOH (Mca: 7-methoxy-coumarin-2-acetic acid; DpaOH: N-3-(2,4-dinitrophenyl)-
25 L-2,3-diaminopropionyl).

The first fraction collected during the purification of each of the pseudopeptides A to G by HPLC (peak 1 of Figure 2 for pseudopeptide G) is used for these tests.

30

From the inhibition profiles obtained with this substrate, for each pseudopeptide A to G, it is possible to determine the constants $K_i N$ and $K_i C$ by following the procedure described by Dive et al. [8].

5

The inhibition experiments were performed at 25°C, pH 6.8, 50 mM HEPES, 10 mM, $CaCl_2$, 200 mM NaCl.

The results obtained are given in the attached Table 1.

10

The same test with pseudopeptide A not comprising the pseudoproline residue was performed for comparative purposes. The results obtained are also given in Table 1. This pseudopeptide was prepared from the phosphinic block ZPhe[PO(OAd)-CH₂]AlaOH, described by Yiotakis et al. [14], followed by coupling of this block to a Wang resin substituted with the residue Fmoc-Ala.

Study of the inhibitory effects of pseudopeptides A to G on ACE and comparison of their affinity towards the N-terminal site ($K_i N$) and the C-terminal site ($K_i C$) of ACE (Table 1) allow the following conclusions to be drawn.

25 **1°) Pseudopeptides A and B:** As regards the affinity for the two active sites of ACE, the presence of a pseudoproline residue appears to be much less favourable than that of a pseudoalanine residue. On the other hand, the presence of a pseudoproline residue in position P1' of these phosphinic pseudopeptides gives access to selective inhibitors of the C-terminal site of ACE.

30

This result demonstrates the essential role of the pseudoproline residue to control the selectivity of the inhibitors with respect to the C-terminal site of ACE.

5 **2°) Pseudopeptides B, C, D and G:** The modifications of
position P2' with the residues alanine, proline,
arginine and tryptophan demonstrate that the nature of
the side chain in this position is also an essential
factor for selectivity. The presence of the proline
10 residue (pseudopeptide C) generates a powerful but
sparingly selective inhibitor of the N and C sites of
ACE. On the other hand, the presence of a tryptophan
residue makes it possible to obtain an extremely
selective inhibitor (pseudopeptide G) of the C-terminal
15 site of ACE.

3°) Pseudopeptides E and F: The substitution of a
pseudophenylalanine residue in position P1 of the
pseudopeptides with a pseudoalanine or pseudo-homo-
20 phenylalanine residue leads to pseudopeptides that are
less powerful and less selective than pseudopeptide G.
This last result demonstrates a lesser importance of
the position P1 of the inhibitors with respect to the
selectivity.

25

The study of the pseudopeptides A to G makes it
possible to conclude that, in the pseudopeptides of the
invention, each position P1, P1' and P2' contributes
towards the selectivity of the interactions. The
30 presence in pseudopeptide G of the pseudophenylalanine,

pseudoproline and tryptophan residues gives it particularly pronounced selectivity.

**Example 11: Demonstration of the *in vivo* properties of
5 pseudopeptide G**

A study is performed *in vivo* in order to check the capacity of pseudopeptide G to inhibit the conversion of angiotensin I into angiotensin II, and to assess its
10 effects on the cleavage of bradykinin, on the one hand, and of the peptide Ac-SDKP, on the other hand.

1) Study protocol

15 This study is performed on batches of male C57BL6/J mice (Iffa Credo) weighing 20 to 23 g each, each batch comprising 6 mice.

To do this, the mice are anaesthetized by means of an
20 intraperitoneal administration of sodium pentobarbital (Sanofi) at a dose of 80 mg/kg of body weight. The right carotid artery is isolated and a catheter (PE10, 0.28x0.61, A-M Systems, Inc.) is inserted into this artery to allow the withdrawal of blood samples, while
25 another catheter (FEP, 0.12x0.67, Carnegie Medecin) is inserted into the ipsilateral jugular vein to allow the administration of the substances. Throughout the experiment, the body temperature of the mice is maintained at 38°C.

In a first stage, the following are administered to the mice of the same batch, via perfusion over 30 minutes:

- either 50 μ L of an isotonic solution (adjusted to pH 7 containing an amount of pseudopeptide G corresponding to a dose of 0.9, 3, 10 or 30 mg/kg of body weight,
- or 50 μ L of physiological saline,
- or alternatively 50 μ L of a solution containing an amount of perindopril corresponding to a dose of 10 mg/kg of body weight, this substance being a powerful mixed inhibitor of ACE (Servier).

Next, the following is administered as a bolus:

- either a mixture comprising 2 μ g of unlabelled angiotensin I and 21 μ Ci of 3 H-angiotensin I in 50 μ L of isotonic solution,
- or a mixture comprising 2 μ g of unlabelled Ac-SDKP and 17 μ Ci of 3 H-Ac-SDKP in 50 μ L of isotonic solution,
- or a mixture comprising 2 μ g of unlabelled bradykinin and 11 μ Ci of 3 H-bradykinin in 50 μ L of isotonic solution.

Samples of arterial blood, of about 50 μ l, are withdrawn 30, 60 and 90 seconds after injection of the angiotensin I mixture or of the bradykinin mixture to the mice that received these mixtures, and 1, 5, 10 and 15 minutes after the start of injection of the Ac-SDKP mixture to the mice that received the latter mixture. In all cases, the blood is collected in preweighed

polypropylene tubes containing 40 µL of water, 10 µL of 80% TFA and 1 µL of heparin. The exact amounts of blood withdrawn are determined by reweighing the tubes. After addition of 195 µL of distilled water, these tubes are
5 placed in an ice bath for 10 minutes and the samples are centrifuged at 4°C to obtain plasma extracts.

The analysis of these extracts is performed by liquid chromatography using an HPLC system (Perkin Elmer 200)
10 linked to a radioelement detector (Z 500-4 cell, Berthold). The chromatographic separations are performed on a Kromasil C18 column (AIT), by injection of 50 µL of samples and using the following mobile phases and elution gradients:

15

- angiotensin I analysis:

mobile phase:

solvent A: CH₃CN/H₂O/TFA (10/90/0.1)

solvent B: CH₃CN/H₂O/TFA (90/10/0.1)

20

elution gradients:

0-30 min: 0-30% B

30-35 min: 30-100% B

- bradykinin analysis:

mobile phase:

25

solvent A: CH₃CN/H₂O/TFA (10/90/0.1)

solvent B: CH₃CN/H₂O/TFA (90/10/0.1)

elution gradients:

0-30 min: 0-25% B

30-35 min: 25-100% B

30

- Ac-SDKP analysis:

mobile phase:

solvent A: H₂O/TFA (100/0.1)

solvent B: CH₃CN/H₂O/TFA (90/10/0.1)

elution gradients:

0-30 min: 0-30% B

5 30-35 min: 30-100% B

The peaks eluted are identified by comparison of their retention times with those shown by the unlabelled substrates (angiotensin I, bradykinin and Ac-SDKP) and
10 by the expected cleavage products (angiotensin II, BK(1-7) and BK(1-5)).

The assays are performed by integration of the area under the corresponding peaks of the chromatogram. The
15 values thus obtained are normalized as a function of the weight of blood taken from each mouse.

The statistical comparisons of the results are performed via the non-parametric Mann-Whitney U test
20 (Statview 5 software).

2) Results

Figure 3 shows, in the form of a bar chart, the mean ±
25 SD values of the angiotensin II/angiotensin I ratio obtained in the case of the mice that received 0.9, 3, 10 and 30 mg of pseudopeptide G per kg of body weight (bars 0.9, 3, 10 and 30, respectively) and also the mean ± SD value obtained for this same ratio in the
30 control mice (bar T), i.e. the mice that received 50 µl of physiological saline. In this figure,

* corresponds to $p < 0.05$, while ** corresponds to $p < 0.01$, by comparison to the control mice.

Figure 3 demonstrates the fact that pseudopeptide G is capable of inhibiting *in vivo* the cleavage of angiotensin I to angiotensin II and that this inhibition is dose-dependent. Thus, the angiotensin II/angiotensin I ratio measured in the control mice is reduced by 50% in the case of the mice treated with 0.9 mg of pseudopeptide G per kg of body weight, and by 90% in the case of the mice treated with 30 mg of pseudopeptide G per kg of body weight.

Figure 4 illustrates, in the form of a bar chart, the mean \pm SD percentages of protection of bradykinin obtained in the case of the mice that received 10 mg of pseudopeptide G per kg of body weight (bar G) and in the case of the control mice (bar T), i.e., in this case, the mice that received 10 mg of perindopril per kg of body weight. In this figure, ** corresponds to $p < 0.01$ by comparison to the control mice.

As may be seen in Figure 4, with the average degree of protection of bradykinin with perindopril arbitrarily set at 100%, this degree of protection is only 9.2% in the case of the mice which received 10 mg of pseudopeptide G per kg of body weight. This pseudopeptide thus appears to afford only a very moderate prevention of cleavage of bradykinin *in vivo*.

Figure 5 shows, also in the form of a bar chart, the mean \pm SD blood concentrations (expressed in pmol/g of blood) of labelled exogenous Ac-SDKP peptide obtained in the case of the mice that received 10 mg of pseudopeptide G per kg of body weight (bar G) and in the case of the control mice (bar T), i.e. the mice that received 50 μ l of physiological saline. In this figure, ** corresponds to $p < 0.01$ by comparison to the control mice.

10

Figure 5 also shows, for comparative purposes, the mean \pm SD blood concentration of labelled exogenous Ac-SDKP peptide in the case of the mice treated with 10 mg of the pseudopeptide RXP407 (described as being a selective inhibitor of the N-terminal site of ACE in references [7] and [8]) per kg of body weight (bar R) and following an identical operating protocol.

15





As shown by Figure 5, pseudopeptide G, at a dose of 10 mg/kg of body weight, appears to have no significant effect on the cleavage of the peptide Ac-SDKP, whereas RXP407 increases the plasmatic content of this peptide 16-fold relative to that observed in the case of the control mice.




25

Thus, *in vivo*, pseudopeptide G very efficiently inhibits the conversion of angiotensin I to angiotensin II, without veritably preventing the degradation of bradykinin, and-even less so that of the peptide Ac-SDKP.

30

Table 1

Formulae	Pseudo-peptide	Ki N nM(10^{-9} M)	Ki C nM(10^{-9} M)
	A	0,8	0,8
	B	450	20
	C	60	4
	D	200	9

	E	8000	60
	F	8000	60
	G	10000	3

References

- [1] Dzan V.J., 2001, *Hypertension* 37, 1047-1052.
- 5 [2] Linz W., Wiemer G., Gohlke P., Unger T., and Scholkens B.A., 1995, *Pharmacol. Rev.* 47(1), 25-49.
- [3] Soubrier F., Alhenc-Gelas F., Hubert C., Allegrini J., John M., Tregear G., and Corvol P., 1988, *Proc. Natl. Acad. Sci. USA* 85(24), 9386-90.
- 10 [4] Wei L., Alhenc-Gelas F., Corvol P., and Clauser E., 1991, *J. Biol. Chem.* 266(14), 9002-8.
- [5] Jaspard E., Wei L., and Alhenc-Gelas F. (1993) *J. Biol. Chem.* 268(13), 9496-503.
- 15 [6] Azizi M., Rousseau A., Ezan E., Guyene T.T., Michelet S., Grognet J.M., Lenfant M., Corvol P., and Menard J., 1996, *J. Clin. Invest.* 97(3), 839-44.
- [7] WO-A-00/01706.
- 20 [8] Dive V., Cotton J., Yiotakis A., Michaud A., Vassiliou S., Jiracek J., Vazeux G., Chauvet M.T., Cuniasse P., and Corvol P. (1991) *Proc. Natl. Acad. Sci. USA* 96(8), 4330-5.
- [9] Junot C., Gonzales M.F., Ezan E., Cotton J., Vazeux G., Michaud A., Azizi M., Vassiliou S., Yiotakis A., Corvol P., and Dive V., 2001, *J. Pharmacol. Exp. Ther.* 297(2), 606-11.
- 25 [10] FR-A-2 676 059.
- [11] EP-A-0 725 075.
- 30 [12] Jiracek J., Yiotakis A., Vincent B., Lecoq A., Nicolaou A., Checler F., and Dive V., Development of highly potent and selective phosphinic peptide

- inhibitors of zinc endopeptidase 24-15 using combinatorial chemistry., 1995, *J. Biol. Chem.* 270(37): 21701-6.
- 5 **[13]** Jiracek J., Yiotakis A., Vincent B., Checlet F. and Dive V., Development of the first potent and selective inhibitor of the zinc endopeptidase neurolysin using a systematic approach based on combinatorial chemistry of phosphinic peptides, 1996, *J. Biol. Chem.* 271(32): 19606-11.
- 10 **[14]** Yiotakis A., Vassiliou S., Jiracek J., and Dive V., Protection of the hydroxy-phosphinyl function of phosphinic dipeptides by adamantyl. Application to the solid-phase synthesis of phosphinic peptides, 1996, *J. Org. Chem.* 61: 6601-6605.
- 15 **[15]** Vassiliou S., Mucha A., Cuniasse P., Georgiadis D., Lucet-Levannier K., Beau F., Kannan R., Murphy G., Knauper V., Rio MC., Basset P., Yiotakis A., and Dive V., Phosphinic pseudo-tripeptides as potent inhibitors of matrix metalloproteinases: a structure-activity study, 1999, *J. Med. Chem.* 42(14): 2610-20.
- 20 **[16]** Georgiadis D., Vazeux G., Llorens-Cortes C., Yiotakis A., and Dive V., Potent and selective inhibition of zinc aminopeptidase A (EC 3.4.11.7, APA) by glutamyl aminophosphinic peptides: importance of glutamyl aminophosphinic residue in the P1 position, 2000, *Biochemistry* 39(5): 1152-5.
- 25 **[17]** Protective groups in Organic Synthesis, Second Edition, T.W. Green and P.G.M. Wuts, John Wiley & Sons, Inc., 309-315.
- 30

- [18] Baylis E.K., Campbell C.D., and Dingwall J.G., 1984, *J. Chem. Soc. Perkin. Trans. I*, 2845.
- [19] Villieras J., Rambaud W.H., and Graff M., 1986, *Synth. Commun.* 16, 149.
- 5 [20] Chen H., Noble F., Roques P., and Fournie-Zaluski M.C., 2001, *J. Med. Chem.* 44, 3523-3530.